

# Geographic variation in fatty acid composition and food source of the commercial clam (*Venerupis decussata*, Linnaeus, 1758), from the Tunisian Coast: Trophic links

S. Bejaoui<sup>a,✉</sup>, D. Boussoufa<sup>a,¥</sup>, K. Telahigue<sup>a,¥</sup>, I. Chetoui<sup>a</sup>, F. Ghribi<sup>a</sup>, I. Rabeh<sup>a</sup> and M. El Cafsi<sup>a</sup>

<sup>a</sup>Unit of Physiology and Aquatic Environment, Biology Department, Faculty of Science of Tunis, University of Tunis El Manar, 2092, Tunis, Tunisia.

¥ These authors contribute equally to this work

✉ Corresponding author: [safa.bejaoui@fst.utm.tn](mailto:safa.bejaoui@fst.utm.tn)

Submitted: 25 May 2018; Accepted: 26 July 2018

**SUMMARY:** Lake and coastal Tunisian areas are rich biodiversity habitats, although little information is available about the distribution of food sources for the inhabitant species. In this study, a fatty acid analysis was used to study the trophic ecology of *Venerupis decussata* communities from 10 sites located along the Tunisian Coast. The richest population in fatty acids was found in S4 followed by S5 and S8, while that of S1, S3 and S10 were the least rich. Results from multivariate analysis confirmed the ecological position of the studied population based on their fatty acid composition. Our results divided the ten studied populations into three similar groups according to their ecological and geographical positions in relation to environmental parameters and food and trophic links. A principal component analysis revealed that diatoms and dinoflagellates were the predominate diets in all the sampling stations. Bacteria and urban discharge dominated the dietary source of clams from S10 and S9. Zooplankton were the preferred diet of *V. decussata* harvested from the two S2 and S3 lakes; although green algae, phytoplankton and detritus were absent from the dietary source of the two previous populations. Despite spatial differences, clams from the north and the south could be easily distinguished from each other, which indicates the utility of this method in the dietary analysis of different food chain links. This study proves that geographic, ecologic and abiotic factors as well as their mutual interaction should be properly investigated in studies focusing on the trophic chains of aquatic ecosystems.

**KEYWORDS:** Estuary and coastal lagoons; Fatty acid composition; Multivariate analysis; Trophic links; Tunisian waters; *Venerupis decussata*

**RESUMEN:** Variación geográfica en la composición de ácidos grasos y fuente de alimento de la almeja comercial (*Venerupis decussata*, Linnaeus, 1758), de las costas tunecinas: cadenas tróficas. Los lagos y las zonas costeras de Túnez son ricos en hábitats de biodiversidad. Sin embargo, hay poca información disponible sobre la distribución de las fuentes de alimentos para las especies residentes. En este trabajo se utilizó el análisis de ácidos grasos para estudiar la ecología trófica de las comunidades de *Venerupis decussata* de 10 localizaciones a lo largo de las costas tunecinas. La población más rica en ácidos grasos se encontró en S4 seguida de S5 y S8, mientras que la de S1, S3 y S10 fueron las menos ricas. Los resultados del análisis multivariante confirmaron la posición ecológica de la población estudiada en función de su composición de ácidos grasos. Nuestros resultados dividieron las diez poblaciones estudiadas en tres grupos similares según sus posiciones ecológicas y geográficas en relación con los parámetros ambientales y con los enlaces tróficos y alimentarios. El análisis de los componentes principales reveló que las dietas predominantes eran las diatomeas y los dinoflagelados en todas las estaciones de muestreo. Mientras que, las bacterias y la descarga urbana dominaron la fuente dietética de almejas de S10 y S9. Sin embargo, el zooplancton fue la dieta preferida de *V. decussata* cosechada de los dos lagos S2 y S3; las algas verdes, el fitoplancton y los detritos estaban ausentes en la fuente dietética de las dos poblaciones anteriores. A pesar de las diferencias espaciales, las almejas del norte y del sur se pueden distinguir fácilmente entre sí, lo que indica la utilidad de este método en el análisis dietético de los diferentes enlaces de la cadena alimentaria. Este estudio demuestra que los factores geográficos, ecológicos y abióticos, así como su interacción mutua deben investigarse adecuadamente en estudios centrados en las cadenas tróficas de los ecosistemas acuáticos.

**PALABRAS CLAVE:** Aguas tunecinas; Análisis multivariable; Composición de ácidos grasos; Enlaces tróficos; Estuario y lagunas costeras; *Venerupis decussata*

**ORCID ID:** Bejoui S <https://orcid.org/0000-0002-7946-2763>, Boussefa D <https://orcid.org/0000-0003-4316-1616>, Telahigue K <https://orcid.org/0000-0001-8841-9911>, Chetoui I <https://orcid.org/0000-0002-2259-5397>, Ghribi F <https://orcid.org/0000-0001-9350-7510>, Rabeh I <https://orcid.org/0000-0002-0307-473X>, El Cafsi M <https://orcid.org/0000-0002-9771-1110>

**Citation/Cómo citar este artículo:** Bejoui S, Boussefa D, Telahigue K, Chetoui I, Ghribi F, Rabeh I, El Cafsi M. 2019. Geographic variation in fatty acid composition and food source of the commercial clam (*Venerupis decussata*, Linnaeus, 1758), from the Tunisian Coast: Trophic links. *Grasas Aceites* 70 (1), e289. <https://doi.org/10.3989/gya.0580181>

**Copyright:** ©2019 CSIC. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License.

**ABBREVIATIONS:** *Venerupis decussata* (*V. decussata*); Polyunsaturated fatty acids (PUFA); Monounsaturated fatty acids (MUFA); Saturated fatty acids (SFA); Total fatty acids (TFA); Eicosapentaenoic (EPA); Docosahexaenoic (DHA); Arachidonic acid (ARA); Temperature (T °C); Salinity (S psu); Chlorophyll a (Ch a); Suspended matter (SPM).

## 1. INTRODUCTION

The littoral of Tunisia covers 1300 Km; it possesses an important particular biodiversity and is considered the richest ecosystem in the Mediterranean Sea (Hattour and Ben Mustapha, 2013). This ecosystem is situated in the most productive area in the Mediterranean Sea, and its high productivity is maintained by a great level of nutrients in both sediment and water column. These transition zones between land and sea can offer unique ecosystem services ranging from trapping contaminants in their sediments to providing nursery areas for marine species and feeding grounds for migratory birds (McLusky and Elliot, 2004). All these specificities provide the littoral of Tunisia with a particular marine and coastal biodiversity.

In fact, the study of the biological functions of marine organisms, as well as seafood quality/quantity, in relation to the surrounding environment, has always been the matter of investigation in different scientific areas such as environmental ecotoxicology and food chemistry (Albergamo *et al.*, 2016; Gomes *et al.*, 2017). However, to comprehend how the stability of a marine ecosystem tolerates it, an analysis of the feeding associations among organisms, especially filter species like bivalves, is very important. In fact, FA analyses have been used extensively to study trophic relationships and as important determinants of ecosystem health and stability in marine food webs (Dalsgaard *et al.*, 2003; Budge *et al.*, 2006).

Bivalves, especially, *V. decussata*, are present at high densities along the sandy beaches of the Tunisian Coast, and they are one of the most important commercial species. This species is considered a potential candidate for shellfish farming in this area. Like others bivalves (Pagano *et al.*, 2016) *V. decussata* are filter-feeders, and consume the phytoplankton suspended in the water column (Hamida *et al.*, 2004). Their ecological and biological preferences have been studied extensively along the Mediterranean Coast (Hamida *et al.*, 2004; Bejaoui *et al.*, 2017). Notwithstanding their position, the nutritional resources and trophic associations of these plentiful groups are inadequately studied. Recent studies analyzing the trophic relationships in marine food chains have been carried on the FA signature to indicate their distribution and preference (Parrish *et al.*, 2015). However, no data about feeding strategies or trophic links of bivalves taken from the Tunisian waters have been reported. In contrast, other international research based on fatty acid composition has been widely used to determine the trophic link between benthic bivalves and primary producers (Irisarri *et al.*, 2014; Kharlamenko *et al.*, 2015). Other studies carried out on spatial and environment changes in food composition have been reported in several aquatic bivalves harvested from

different localities, such as *Mytilus galloprovincialis* and *Crassostrea gigas* from the Thau Lagoon (Pernet *et al.*, 2012) and *Glauconome chinensis*, *Sinonovacula constricta* from Yangtze Estuarine Intertidal Marsh (Wang *et al.*, 2015).

The current study constitutes a novel investigation on the relationship between the fatty acid composition and the trophic marker of the European clam *V. decussata* and its geographic repartition in the inshore Tunisian area.

The objectives of this study were, then, to answer several questions:

- Does the fatty acid composition of this species vary among geographically different locations?
- Are there any similarities in the quality of the diet supplied by different locations to the same species *V. decussata*?
- Does *V. decussata* have a selective choice for its food?

## 2. MATERIALS AND METHODS

### 2.1. Sampling procedures

Specimens of *V. decussata* (mean total weight:  $8.19 \pm 0.717$  g; mean total length:  $38.25 \pm 5.45$  mm;  $n = 50$  per sites) were sampled by hand fishing on foot or by scuba diving, at ten different commercial fishing sites along the Tunisian Coast (Figure 1 and Table 1), during the winter season, 2015. The sampling sites were: Bizerte Lagoon (S1); North Lake (S2); South Lake (S3); Louza (S4); Zabboua (S5); Bousaid (S6); Bounglow (S7); Maoumma (S8); Zarrat (S9) and Boughrara Lagoon (S10). The harvested samples were immediately transported to the laboratory in ice boxes. The clams were then kept in filtered sea water flow-through aquaria for at least 24 h to depurate their gut before being dissected. This procedure ensured the overall expulsion of ingested food accumulated in the mid-gut (Boussoufa *et al.*, 2011). For each site, the soft tissues of the clams ( $35.5 \pm 3.1$  mm anteroposterior shell length) were placed in  $-30$  °C for fatty acid analysis. Temperature, salinity and pH were measured in situ with a thermometer, salinity-conductivity (model: WTW. 800.645.5999) and pH meter (model: WTW.LF.325), respectively. In the laboratory, suspended matter and chlorophyll a were determined according to the Aminot and Chaussepied (1983) methods. In fact, the suspended matter content was obtained by the filtration of 500 mL of water sample, added at the end of the filtration with 20 mL of format ammonium in order to remove all traces of salt on the filter. Nucleopore Track-Etch Whatmann filter paper (porosity  $0.45$   $\mu$ m) was weighed (dry weight of the front filter filtration) after placing in an oven at  $60$  °C for 24 hours. After filtering, the water samples from each station were stored for at




FIGURE 1. Sampling stations of *V. decussata* along the Tunisian coast.

Bizerte Lagoon (S1), Chekly (S2), Baie (S3), Louza (S4), Zabbousa (S5), Boussaid (S6), Bunglow (S7), Maoumma (S8), Zarrat (S9) and Boughrara Lagoon (S10).

least 48 hours in an oven at 60 °C to dry. Then the filters were weighed with a precision balance and the dry weight of suspended solids was obtained by subtracting the dry weight of the filter after filtration from the empty one. Finally, the concentration of suspended matter was obtained by dividing by the volume of filtered water.

## 2.2. Analysis of the fatty acid composition

The lipid analysis was carried out on the whole soft body homogenate, extracted with chloroform/methanol (2v/1v), containing 0.01% butylated hydroxytoluene (BHT) as an antioxidant, according to the method of Folch *et al.*, (1957). The lipid content was determined by the gravimetric method and expressed as percent wet weight (WW). The fatty acids (FA) from the total lipids were *trans*-methylated with a solution of sodium methylate

and concentrated sulfuric acid solution in methanol (2%), according to Cecchi *et al.*, (1982). The mixture was centrifuged at 4000 g for 10 min. The supernatant containing the total fatty acid methyl esters (FAME) was injected into a gas chromatograph equipped with a flame ionization detector and a 30 m capillary column of flexible silica 250 µm in diameter and 0.25 µm film thickness (Supelco, PUFA -3). A temperature injector injector of 275 °C was used, operating in a solvent elimination mode. Nitrogen was the carrier gas. Identification of FAMES was based on the comparison of their retention times with those of a mixture of methyl esters (SUPELCO PUFA-3), authentic standards (C4 C24 by SUPELCO) and to a well-characterized fish oil (Mehaden oil by SUPELCO). Fatty acid peaks were integrated and analyzed using HP chemstation software. The relative amount of each FA was expressed as % of total fatty acids (%TFA).

TABLE 1. Characteristics of the sampling locations along the Tunisian waters (Center Mediterranean Sea).

Regions	Sampling Stations	Geographical position	Characteristics
northern locations	S1 (Lagoon of Bizerte)	N37°13'25.45"E9°55'31,30"	<ul style="list-style-type: none"> <li>Connected with three hydric components: the maritime contribution, Oueds (Mrazig, Graa, Ben Hassna) and the exchanges by the intermeddle channel (tinja) with Ichkul Lake.</li> <li>This lagoon is rich in suspension material, chlorophyll a, and nutrient salts. However, there is depletion of phytoplankton and zooplankton levels.</li> <li>This lagoon receives daily anthropogenic disturbances from various origins: e.g. domestic, industrial and agricultural wastes released by the neighboring agglomerations.</li> </ul>
	Lagoon of Tunisia S2 (Chekly)	N36°49'4.31"- E10°13'5.94"	<ul style="list-style-type: none"> <li>This lagoon undergoes a significant degradation of waters due to the urban wastewater discharge.</li> </ul> <p>This site is located in the northern part of this lagoon. It is characterized by sandy sediments with some zones of sticky mud rich in seaweeds.</p>
	S3 (Baie)	N36°47'12,40"-E10°13'12,88"	<ul style="list-style-type: none"> <li>This lagoon was eutrophic.</li> <li>It is connected with the Gulf of Tunisia by a channel of Kheireddine</li> </ul> <p>This lagoon is rich in nutrient salts and has an important resource.</p> <p>This site is situated in front of the mouth of the Oued Sidi Daoued. It is characterized by a black sticky mud with a strong smell of hydrogen sulfide (H<sub>2</sub>S) and very rich in seaweeds with some fragments of shells.</p>
central locations	S4 (Louza)	N35°0'44.10" – E11°0'47.48"	<ul style="list-style-type: none"> <li>The site of Louza has been considered as a reference site in monitoring programs along the Tunisian Coast.</li> </ul>
southern locations	S5 (Zabbousa)	N34°24'32.09" – E10°22'14.56"	<ul style="list-style-type: none"> <li>The Gulf of Gabes is an ecosystem which has taken a favored place in the center of the Mediterranean Sea. This gulf is characterized by a strong tide and swell, heterogeneous sediment, sub-desert climate and specific diversity.</li> <li>This gulf is rich in suspension mater, chlorophyll a and nutrient salts.</li> <li>Connected with the Mediterranean Sea by two channels (Ajim and El Kantara).</li> <li>Receives an important land discharges via six Oueds.</li> <li>The sedimentation is sandy, muddy and mixed.</li> <li>This ecosystem is fragile and it is characterized by irregular hydrodynamics which cause massive eutrophication</li> <li>This lagoon is polluted. It receives important discharges daily from the aquaculture, industrial and urban activities.</li> </ul>
	S6 (Boussaid)	N34°12'5.10" – E10°3'8.86"	
	S7 (Bunglow)	N33°51'14,49"- E10°9'13.17"	
	S8 (Maoumma)	N33°48'10,75"-E10°12'17.61"	
	S9 (Zarrat)	N33°39'16.66" - E10°28'25.08"	
	S10 (Lagoon of Boughrara)	N33°32'34.20" -E10°40'56.37"	

### 2.3. Fatty acid trophic marker

Fatty acid ratios were determined and applied to assess broad clams' food web from the Tunisian Coast. Among them, diatom feeders:  $\Sigma 16$ , EPA (Dalsgaard *et al.*, 2003); C16:1/C16:0 (Prato *et al.*, 2010) and C18:1+C16:1n-7 was reported by Auel *et al.*, (2002). Also, C20:1+C22:1 was proposed to reflect the organisms' diet by zooplankton (Parrish *et al.*, 2015). The high proportion of C15:0+C17:0+C18:1 denotes the presence of bacteria in the *V. decussata* diet (Budge *et al.*, 2001). A green algae diet was determined by the presence

of C16:3; C18:3; C20:4 (Leveillé *et al.*, 1997). The C16:0, C18:4n-3 and high DHA/EPA ratio (>1) reflects the proportion of dino flagellates (Budge *et al.*, 2001; Dalsgaard *et al.*, 2003). PUFAs 22 and HUFAs n-3/n-6 were described by Desvillettes *et al.*, (1997a) and Rocchetta *et al.*, (2014) characterized the detritus feeders. Cyanophytes and phytoplankton were also determined by C16:0, C16:1n-7, C16:4n-3, C18:0 and C18:2n-6 (Li and watanabe, 2001; Irisarri *et al.*, 2014). One other fatty acid trophic marker such as C18:1+C18:2n-6 was evaluated to esteemed urban discharge (Sakdullah and Tsuchiya, 2009).



## 2.4. Statistical analysis

All analyses were carried out in ten replicates except fatty acids (triplicate). The results were expressed as mean values  $\pm$  standard deviation (SD). A multivariate statistical analysis was performed using Primer 7 software analysis. Similarity, FA composition was checked by the Bray-Curtis test, without a pre-treatment. The impact of individual FAs on the similarities and dissimilarities among sample sites were analysed using the SIMPER analysis. FA groups were identified using the factorial correspondence analysis (FCA). Trophic markers were analysed through the principal component analysis (PCA) using the “R” software version 2.15.3 (R Core Team, 2012). We further used non-metric multidimensional scaling (NMDS), calculated and plotted in R Vegan library as a multivariate visualization tool for the whole data set. Data were analysed for normality variance homogeneity through the Shapiro test (Sokal and Rohlf, 1987). Post hoc tests of one-way ANOVA followed by Tukey HSD test and non-parametric Kruskal-Wallis test for samples with unequal variances were performed so as to investigate significant differences among biochemical parameters. Statistical analyses were performed for a significance level at 0.05 using Statistica (Version 8).

## 3. RESULTS

### 3.1. Abiotic factors

The physicochemical parameters of the ten sampling sites are summarized in Figure 2. During winter, the temperature ( $T$  °C) and salinity ( $S$  psu) ranged from 13 to 17 °C and from 29 to 35 psu, respectively, with significant differences among sites ( $p < 0.05$ ). Chlorophyll a (Chl a) and suspended material (SPM) followed the same fluctuation within stations with significantly higher values in S2 (northern station: 275.71 mg/L; 1.49  $\mu\text{g}/\text{cm}^3$  for SPM and Chl a; respectively) and S10 (southern station: 306.26 mg/L; 1.40  $\mu\text{g}/\text{cm}^3$ , for SPM and Chl a; respectively) ( $p < 0.05$ ) and the lower one in S6 and S7 (southern stations) in the order of 0.82 and 0.69  $\mu\text{g}/\text{cm}^3$  for SPM and 143.74 and 150.99 mg/L for Chl a ( $p < 0.01$ ).

### 3.2. Fatty acid composition

In general, PUFAs were present in the highest percentages in all the studied sites, with maximum values recorded in clams from S3 (40%) and S4 (60%) (Table 2). Among them, linoleic (C18:2n-6) (>35%), docosahexaenoic (DHA) (>13%), eicosapentaenoic (EPA) (>6%) and arachidonic (ARA) (6%) acids had a high range of relative percentage and varied significantly among the sampling stations ( $p < 0.01$ ). Nevertheless, DHA varied significantly in




FIGURE 2. Abiotic parameters of the ten sampled sites along the Tunisian Coast.

Bizerte Lagoon (S1), Chekly (S2), Baie (S3), Louza (S4), Zabbousa (S5), Boussaid (S6), Bunglow (S7), Maoumma (S8), Zarrat (S9) and Boughrara Lagoon (S10).

Suspended mater (SPM);  $T$  °C (Temperature); Spsu (Salinity); Chlorophyll a (Chl a).

Values are expressed as means  $\pm$  SD of three replicates.

Significant difference is given by asterisk at 0.05: \* $p < 0.05$  and \*\* $p < 0.01$  using the ANOVA test (Tukey HSD).

*V. decussata* from the different stations and was the highest in S2 with 13% of TFA. The lowest percentages were registered in clams from S9 and S10 (<2%) ( $p < 0.001$ ). Dissimilar variations were recorded for C18:2n-6 with high percentages in specimens from S2 (1.24%) as compared to those from S9 and S10 (>26%). However, similar trends were observed for the EPA and ARA acid percentages, which presented the highest levels in *V. decussata* from S3 (6.59% and 2.90%, respectively) and the lowest levels in the clams from S9 (0.39% and 0.81%, respectively) and S10 (0.65% and 0.59%, respectively). The notable

TABLE 2. Fatty acid composition of *V. decussata* sampled from ten sites along the Tunisian Coastline, expressed as percentage of total fatty acids.

Fatty Acids	S1 (bizerte Lagoon)	S2 (Chekly)	S3 (Baie)	S4 (Louza)	S5 (Zabbousa)	S6 (Boussaid)	S7 (Bunglow)	S8 (Maoumma)	S9 (Zarrat)	S10 (Boughrara)	F	p
<i>C14:0</i>	0.884±0.067	4.579±1.578	5.063±0.436	1.398±0.510	2.575±1.139	2.712±0.834	3.051±1.381	2.486±0.274	1.358±0.680	1.609±0.400	7.9506	<0.001 (***)
<i>C15:0</i>	1.826±0.080	0.256±0.081	1.392±0.266	1.364±0.726	1.614±0.330	1.449±0.862	1.415±0.643	0.724±0.100	0.417±0.257	0.659±0.419	4.8074	<0.001 (***)
<i>C16:0</i>	23.566±2.069	17.601±1.973	21.585±3.337	11.450±5.542	9.217±7.938	12.005±6.345	18.587±9.179	11.459±0.988	15.930±0.987	19.071±1.877	3.0830	0.008 (**)
<i>C18:0</i>	5.358±0.040	8.505±1.087	7.555±0.574	6.707±3.096	9.375±1.182	8.401±4.036	8.403±1.720	7.049±0.095	4.748±0.399	5.004±0.172	2.3828	0.03 (*)
<i>C20:0</i>	2.263±0.095	0.616±0.239	0.186±0.129	0.807±0.734	0.265±0.148	1.843±1.829	1.874±0.802	2.210±0.013	0.680±0.090	1.418±0.526	3.7469	0.002 (**)
<i>C22:0</i>	0.739±0.675	0.114±0.064	0.129±0.070	2.516±2.297	1.376±0.690	0.506±0.515	0.030±0.025	0.090±0.004	0.031±0.029	0.116±0.100	3.9819	0.0016 (**)
<b>Σ Saturates</b>	34.637±2.527	31.674±4.030	35.913±2.844	24.245±8.670	24.425±7.501	26.920±8.044	33.351±9.351	24.021±0.918	23.166±2.188	27.879±2.785	3.897	0.02 (*)
<i>C14:1</i>	1.681±0.663	0.750±0.243	0.525±0.292	1.241±0.329	1.204±0.584	1.594±1.054	0.872±0.419	1.477±0.119	0.508±0.493	0.336±0.176	2.7709	0.015 (*)
<i>C15:1</i>	1.349±0.609	1.022±0.266	0.810±0.406	2.223±1.811	1.655±0.616	1.568±0.283	1.330±1.355	0.591±0.074	0.308±0.170	0.447±0.247	1.9317	NS
<i>C16:1n-11</i>	8.197±4.403	4.079±1.569	5.853±1.497	4.082±2.334	1.905±1.116	2.090±0.931	2.761±2.487	2.313±0.088	0.645±0.237	0.849±0.243	5.6274	<0.001 (***)
<i>C16:1n-9</i>	2.041±0.168	1.259±0.184	1.276±0.260	2.534±1.317	2.617±0.793	5.344±3.970	3.172±1.700	2.101±0.237	0.968±0.449	0.740±0.263	3.0049	0.009 (**)
<i>C16:1n-7</i>	2.363±0.607	1.678±0.454	1.736±1.002	1.376±0.337	2.976±1.428	2.932±1.800	2.613±0.763	4.603±0.304	1.429±0.436	2.561±0.944	2.6837	0.018 (*)
<i>C18:1</i>	5.438±1.806	7.932±1.104	7.780±1.927	2.080±2.930	4.594±4.373	3.618±3.764	5.755±2.398	4.516±1.346	21.160±1.816	24.245±3.396	22.7939	<0.001 (***)
<i>C20:1n-9</i>	2.129±0.673	7.185±1.663	4.899±0.989	1.062±0.320	1.977±0.695	1.558±0.708	1.221±0.155	1.257±0.158	0.818±0.337	0.552±0.118	31.8972	<0.001 (***)
<i>C22:1n-9</i>	0.202±0.035	0.275±0.120	0.243±0.186	0.491±0.209	0.548±0.427	0.349±0.294	0.666±0.214	0.564±0.291	0.332±0.251	0.203±0.100	1.7958	NS
<b>Σ Monoenes</b>	23.404±5.635	24.183±3.305	23.126±3.220	15.093±2.414	17.481±4.333	19.058±5.616	17.841±3.972	17.425±1.544	26.169±0.144	29.936±2.303	20.368	0.03 (*)
<i>C16:2</i>	2.014±0.574	0.460±0.218	0.441±0.140	3.250±1.737	4.389±1.067	2.507±0.864	2.935±2.204	1.396±0.201	1.107±0.320	0.888±0.427	6.7261	<0.001 (***)
<i>C16:3</i>	1.920±0.703	1.482±0.086	1.878±0.849	2.085±0.620	5.040±4.197	1.393±1.307	2.530±0.962	1.002±0.154	0.701±0.518	0.515±0.277	2.6467	0.019 (*)
<i>C16:4</i>	7.910±1.617	0.512±0.161	0.477±0.190	4.159±2.907	8.287±2.391	6.283±4.326	4.116±2.848	6.157±0.592	1.523±0.728	1.260±0.331	6.7421	<0.001 (***)
<i>C18:2n-6</i>	1.983±0.021	1.249±0.290	1.765±0.849	2.858±1.318	2.996±0.439	6.394±3.609	3.507±0.875	3.083±0.400	35.268±4.943	26.521±3.229	103.6171	<0.001 (***)
<i>C18:3n-6</i>	0.268±0.122	0.592±0.376	0.562±0.383	0.514±0.433	1.457±0.971	1.325±0.958	2.649±1.662	5.104±0.162	1.154±0.898	0.386±0.127	8.0070	<0.001 (***)
<i>C18:3n-3</i>	2.121±0.256	1.413±0.792	1.584±1.059	1.751±0.518	1.633±0.270	1.593±0.623	0.836±0.345	0.720±0.014	3.657±0.494	2.810±0.450	7.0754	<0.001 (***)
<i>C18:4n-3</i>	0.560±0.320	1.613±0.309	2.311±1.296	1.261±0.213	0.561±0.549	1.504±0.514	0.994±0.430	0.654±0.010	0.564±0.116	0.719±0.360	4.8129	<0.001 (***)
<i>C20:2n-6</i>	0.394±0.147	1.663±0.205	1.382±1.004	6.354±6.716	1.980±0.418	8.839±7.220	11.057±7.289	16.722±0.689	0.536±0.067	0.500±0.270	5.0775	<0.001 (***)
<i>C20:3n-6</i>	0.259±0.098	2.161±0.563	1.762±0.847	1.367±1.170	1.056±0.519	3.769±3.799	1.410±1.086	3.923±0.001	0.778±0.478	2.317±0.840	2.0057	NS
<i>C20:4n-6 ARA</i>	1.829±1.607	2.261±0.551	2.909±1.220	1.450±1.122	2.140±1.456	1.250±0.855	2.630±1.084	2.806±0.301	0.813±0.350	0.596±0.098	2.2415	0.04 (*)
<i>C20:3n-3</i>	0.243±0.009	0.085±0.048	0.136±0.103	0.353±0.552	0.085±0.080	0.782±1.071	0.252±0.133	0.758±0.007	0.114±0.022	0.408±0.198	1.3506	NS
<i>C20:4n-3</i>	3.757±0.238	0.285±0.049	0.344±0.086	1.461±1.566	0.181±0.040	0.129±0.083	0.152±0.066	0.196±0.029	0.026±0.013	0.021±0.005	14.6747	<0.001 (***)
<i>C20:5n-3 EPA</i>	0.599±0.520	4.043±0.644	6.594±3.649	1.939±1.154	2.218±1.155	2.426±1.408	2.184±1.313	2.557±0.240	0.397±0.168	0.658±0.014	6.0579	<0.001 (***)
<i>C22:2n2j</i>	3.513±1.235	5.458±1.816	3.620±2.329	5.288±1.859	4.240±0.826	2.093±1.753	2.464±1.827	2.913±0.579	0.851±0.371	1.209±0.379	4.2861	<0.001 (***)
<i>C22:2n-6</i>	1.429±0.579	1.189±0.316	1.031±0.481	2.403±1.387	2.123±0.754	1.366±2.116	0.967±0.724	0.879±0.107	0.407±0.193	0.345±0.067	1.6427	NS
<i>C22:5n-6</i>	2.449±0.991	2.453±1.791	1.270±0.589	6.753±4.158	4.420±2.086	3.392±3.050	2.135±1.698	1.636±0.355	0.624±0.449	0.476±0.178	3.2006	0.006 (**)
<i>C22:5n-3</i>	4.133±0.902	4.065±2.519	1.974±0.991	6.767±3.555	5.644±3.862	3.303±3.424	2.106±1.698	1.536±0.146	0.534±0.286	0.632±0.178	2.5629	0.02 (*)
<i>C22:6n-3 DHA</i>	6.567±1.968	13.147±3.630	11.050±2.359	10.637±3.972	9.635±1.062	5.898±3.378	5.873±3.736	6.503±1.403	1.600±0.784	1.912±0.337	7.1977	<0.001 (***)
<b>Σ polyunsaturates</b>	41.958±8.160	44.142±6.553	40.960±2.023	60.660±7.439	58.092±10.921	54.021±11.697	48.807±11.411	58.552±2.463	50.663±2.325	42.183±4.275	123.33	<0.001 (***)
<b>Σ n-3 PUFA</b>	19.833±2.244	23.065±5.691	18.672±2.674	28.987±7.165	22.161±6.465	16.603±0.877	12.351±5.518	12.006±1.937	7.122±1.330	6.982±1.159	80.68	<0.001 (***)
<b>Σ n-6 PUFA</b>	4.336±0.734	6.857±0.658	6.364±1.102	13.499±5.653	9.614±1.393	21.695±2.997	19.592±7.147	29.712±1.142	38.144±4.437	30.070±2.692	67.39	<0.001 (***)

Values are expressed as means ± SD of ten pooled clams for each site and three biological replicates were produced.

Significant difference are given by asterisk at 0.05; \*p < 0.05; \*\*p < 0.01 and \*\*\*p < 0.001 using the ANOVA test (Tukey HSD).

ARA: arachidonic acid, DHA: docosahexaenoic acid, EPA: eicosapentaenoic acid.

amounts of the specific NMID FA ( $\Sigma C22:2i/2j$ ) found in the clams from S2 (5%), S4 (5%) and S5 (4%) were significantly higher than those found in S9 (0.8%) and S10 (1%) ( $p < 0.001$ ). With regard to the relative participation of the saturated (SFA) and monounsaturated (MUFA) fatty acids, both groups presented lower percentage throughout the study as compared to PUFA ( $<35\%$  and  $<29\%$ ). In fact, the maximum amount of SFA was recorded in the clams from S3 with 35%, but, the minimum was obtained from S9 with 23% of TFA ( $p < 0.05$ ). In this context, SFA was dominated by C16:0 and C18:0 acids ranging from 9 to 23% and from 4 to 9% of TFA, respectively ( $p < 0.05$ ). Concerning MUFA, the main proportion was observed in clams from S9 and S10 ( $<24\%$  of TFA). This group was predominated by C18:1, which varied significantly among the studied stations ( $p < 0.001$ ). The SIMPER analysis indicated that n-3 PUFA was the main differentiating series due to its elevated presence in the Bizerte (19%), North Lagoon (23%), South Lagoon (18%), Louza (28%) and Zabbousa (22%) areas. However, the Boughrara Lagoon and the Zarrat areas were characterized

by the lowest percentages of this series ( $p < 0.05$ ). Conversely, the amount of PUFA (n-6) was greater in the Boughrara (S10; 30%) and Zarrat (S9; 38%) stations than in Bizerte (S1) and the north Tunisian lagoons (S2, 4% and S3, 6%, respectively) ( $p < 0.001$ ).

Multivariate statistical analyses are represented in Figure 3 and Table 3. The FCA analysis separated the stations into three groups. The FCA showed that the first three factors contributed to 63.97 % of the total inertia (factor 1:36.75%; factor 2: 14.60% and factor 3: 12.62%). The first one (Group A) regrouped the bivalves sampled from the north coast area (S1, S2 and S3) at close to 96.23%, characterized by a higher contribution of C16:0, DHA, C14:0, C22:2i/2j, EPA and C16:1-n11 fatty acids at 63.14%, 59.17%, 44.21%, 43.53%, 43.53% and 41.42%; respectively. The specimens from S4, S5, S6, S7 and S8 (Group B; close to 92.14%) contained an appreciable amount of C20:2n-6, C22:5n-3, C22:5n-6, C16:4n-3, C22:2n-6, C16:2 and C18:3n-6; and explained 90.62%, 69.44%, 62.70%, 59.90%, 47.41%, 44.47% and 40.48% of group similarity, respectively. Only C18:2n-6 was present in




FIGURE 3. Factorial correspondence analysis (FCA) of the fatty acid composition of *V. decussata* sampled along the Tunisian Coast. N: sites from the north part of Tunisia including Bizerte Lagoon (S1), Chekly (S2) and Baie (S3). C: sites from the central part of Tunisia including Louza (S4), Zabbousa (S5), Boussaid (S6), Bunglow (S7) and Maoumma (S8). S: sites from the south part of Tunisia including Zarrat (S9) and Boughrara Lagoon (S10).

TABLE 3. Similarity and dissimilarity multi-analysis (PRIMER I) of the fatty acid composition of *V. decussata* tissues sampled along ten sites from the Tunisian water.

Variables	Similarity										Dissimilarity		
	Group A			Group B				Group C					
	S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	Groups A vs B	Group A vs C	Group B vs C
Fatty acids													
Cumulative %	96.23%			92.14%				84.73%			74.58%	72.62%	64.62%

higher amounts in Group C than the other groups in the order of 97.58%. This group, characterizing *V. decussata* from S9 and S10, contained a similarity in FA composition of 84.73%. In Table 3, SIMPER provides a quantitative comparison among sites with a high dissimilarity between groups (>50%). The lowest dissimilarity was observed between Groups B and C at 64.62%. However, the highest one was detected between Group A and B at 74.58%.

Non-metric multidimensional scaling was used in this work to better understand the relationship between the lipid compositions of clams and environmental parameters (Figure 4). The total FA composition differed significantly among the sampling sites. The two-dimensional NMDS plot based on the relative presence of different FA and abiotic parameters revealed a clear separation according to their geographical location. The FA composition of the three northern locations appeared particularly close among themselves. The Person correlation analysis revealed significant positive correlations between FA compositions of *V. decussata* collected from the three northern locations and abiotic parameters such as SPM (0.86); S psu (0.99) and Ch a (0.88). A reasonably large amount of variation in the fatty

acid compositions and abiotic parameters occurred within each group. In this context, differences were observed among sampling areas and the measured environment variation, in particular, S psu ( $r = 0.22$ ;  $p < 0.009$ ) and Ch a ( $r = 0.22$ ;  $p < 0.01$ ) were found to cluster separately in the northern locations from the other one. However, T °C ( $r = 0.48$ ;  $p < 0.01$ ) was negatively correlated with the two southern locations ( $-0.87$ ). The other main differentiating variable in those latter locations was linoleic acid (C18:2 $\omega$ 6; 26% and 35% contribution), which was accumulated in high levels in clam tissues from the southern locations. The bi-plot analysis suggested no significant correlation between the FA compositions of the clams collected from the center coast areas and the environmental parameters in this study. However, specimens from the southern region were differentiated from the others groups by C20:2n-6.

### 3.3. Fatty acid trophic markers

The fatty acid trophic markers are summarized in Table 4. Ten food source indicators were determined in our study, dinoflagellates, diatoms, cyanophytes, bacteria, zooplankton, phytoplankton, detritus,




FIGURE 4. Non-metric multidimensional scaling (NMDS) analysis between fatty acid compositions of *V. decussata* and environmental parameters.

N: sites from the north part of Tunisia including Bizerte Lagoon (S1), Chekly (S2) and Baie (S3).

C: sites from the central part of Tunisia including Louza (S4), Zabbousa (S5), Boussaid (S6), Bunglow (S7) and Maoumma (S8).

S: sites from the south part of Tunisia including Zarrat (S9) and Boughrara Lagoon (S10).

SUSPENDED mater (SPM); T °C (Temperature); Spsu (Salinity); Chlorophyll a (Chl a).



TABLE 4. Trophic markers in *V. decussata* sampled among ten sites from the Tunisian water (expressed as percentage of total fatty acids).

Types	FA biomarkers	S1 (bizerte Lagoon)	S2 (Chekty)	S3 (Baie)	S4 (Louza)	S5 (Zabbousa)	S6 (Boussaid)	S7 (Bunglow)	S8 (Maoumma)	S9 (Zarrat)	S10 (Boughrara)	p
<i>Diatoms</i>	$\Sigma 16$ specific	48.013 $\pm$ 3.660	27.074 $\pm$ 3.656	33.250 $\pm$ 4.086	28.776 $\pm$ 6.903	36.816 $\pm$ 1.217	32.325 $\pm$ 7.098	36.164 $\pm$ 8.810	29.033 $\pm$ 0.896	22.305 $\pm$ 3.084	25.888 $\pm$ 2.333	<0.01 (**)
	C20 :5n3	0.599 $\pm$ 0.520	4.043 $\pm$ 0.644	6.594 $\pm$ 3.649	2.014 $\pm$ 1.319	2.657 $\pm$ 0.473	2.426 $\pm$ 1.408	2.184 $\pm$ 1.313	2.557 $\pm$ 0.240	0.397 $\pm$ 0.168	0.658 $\pm$ 0.014	<0.001 (***)
	C16 :1/C16 :0	0.102 $\pm$ 0.036	0.096 $\pm$ 0.028	0.086 $\pm$ 0.060	0.232 $\pm$ 0.292	0.901 $\pm$ 1.072	0.333 $\pm$ 0.243	0.192 $\pm$ 0.162	0.402 $\pm$ 0.008	0.088 $\pm$ 0.023	0.134 $\pm$ 0.048	<0.001 (***)
	C16 :1n7+C18 :1	7.801 $\pm$ 1.203	9.610 $\pm$ 1.241	9.517 $\pm$ 2.754	2.154 $\pm$ 1.019	8.481 $\pm$ 2.765	6.551 $\pm$ 4.509	8.368 $\pm$ 2.966	9.119 $\pm$ 1.651	22.589 $\pm$ 1.521	26.806 $\pm$ 2.827	<0.01 (**)
<i>Green algae</i>	$\Sigma 16$ , 18, 20-n3	7.799 $\pm$ 0.336	3.181 $\pm$ 0.783	3.808 $\pm$ 1.618	5.299 $\pm$ 1.171	7.293 $\pm$ 4.684	2.884 $\pm$ 0.776	3.518 $\pm$ 1.229	1.918 $\pm$ 0.169	4.386 $\pm$ 0.350	3.348 $\pm$ 0.241	<0.05 (*)
	$\Sigma$ PUFAs 18	4.374 $\pm$ 0.155	3.255 $\pm$ 0.570	3.772 $\pm$ 2.155	5.125 $\pm$ 1.791	5.669 $\pm$ 0.629	9.313 $\pm$ 3.405	6.994 $\pm$ 2.045	8.907 $\pm$ 0.576	40.080 $\pm$ 5.204	29.718 $\pm$ 3.581	<0.001 (***)
	C18 :3n3	2.121 $\pm$ 0.256	1.413 $\pm$ 0.792	1.584 $\pm$ 1.059	1.751 $\pm$ 0.518	1.620 $\pm$ 0.300	1.593 $\pm$ 0.623	0.836 $\pm$ 0.345	0.720 $\pm$ 0.014	3.657 $\pm$ 0.494	2.810 $\pm$ 0.450	<0.001 (***)
<i>Dinoflagellates</i>	C16 :0	23.034 $\pm$ 2.620	17.601 $\pm$ 1.973	21.585 $\pm$ 3.337	11.541 $\pm$ 7.827	10.651 $\pm$ 7.958	12.005 $\pm$ 6.345	21.795 $\pm$ 6.615	4.516 $\pm$ 1.346	15.930 $\pm$ 0.987	19.071 $\pm$ 1.877	<0.001 (***)
	C18 :4n3	0.627 $\pm$ 0.421	1.613 $\pm$ 0.309	2.311 $\pm$ 1.296	1.212 $\pm$ 0.263	0.662 $\pm$ 0.548	1.504 $\pm$ 0.514	1.062 $\pm$ 0.464	6.503 $\pm$ 1.403	0.564 $\pm$ 0.116	0.719 $\pm$ 0.360	<0.001 (***)
	C22 :5n-3/C20 :5n-3	7.965 $\pm$ 0.165	3.203 $\pm$ 0.466	2.263 $\pm$ 1.517	4.626 $\pm$ 3.104	3.624 $\pm$ 0.366	2.431 $\pm$ 0.868	1.750 $\pm$ 0.945	2.528 $\pm$ 0.310	3.954 $\pm$ 0.271	2.897 $\pm$ 0.458	<0.05 (*)
<i>Zooplankton</i>	C20 :1+C22 :1	2.332 $\pm$ 0.665	7.460 $\pm$ 1.705	5.143 $\pm$ 1.128	1.554 $\pm$ 0.336	2.669 $\pm$ 0.839	1.908 $\pm$ 0.910	1.888 $\pm$ 0.224	1.822 $\pm$ 0.450	1.150 $\pm$ 0.485	0.755 $\pm$ 0.085	<0.01 (**)
<i>Bacteria</i>	C15 :0+C17 :0+C18 :1n7	7.264 $\pm$ 1.734	8.188 $\pm$ 1.104	9.173 $\pm$ 1.723	2.243 $\pm$ 0.577	7.031 $\pm$ 4.440	5.068 $\pm$ 3.093	7.171 $\pm$ 1.959	5.240 $\pm$ 1.447	21.577 $\pm$ 1.750	24.904 $\pm$ 3.145	<0.01 (**)
<i>Detritus</i>	PUFAs 22-n6	3.878 $\pm$ 1.568	3.642 $\pm$ 1.868	2.302 $\pm$ 1.010	9.157 $\pm$ 5.362	5.788 $\pm$ 2.025	6.932 $\pm$ 2.992	2.788 $\pm$ 1.523	2.515 $\pm$ 0.463	1.032 $\pm$ 0.607	0.821 $\pm$ 0.217	<0.001 (***)
	HUFAs n6+n3	21.664 $\pm$ 4.452	28.905 $\pm$ 5.669	27.187 $\pm$ 0.773	39.489 $\pm$ 9.610	27.522 $\pm$ 3.959	37.656 $\pm$ 5.564	23.991 $\pm$ 4.595	37.521 $\pm$ 1.903	5.833 $\pm$ 2.565	7.393 $\pm$ 1.344	<0.001 (***)
<i>Urban discharges</i>	C18 :1+C18 :2n6	7.422 $\pm$ 1.804	9.181 $\pm$ 1.218	9.546 $\pm$ 1.303	3.626 $\pm$ 1.021	8.298 $\pm$ 4.337	10.865 $\pm$ 4.940	9.262 $\pm$ 1.754	7.599 $\pm$ 1.746	56.428 $\pm$ 6.756	50.766 $\pm$ 4.561	<0.001 (***)
<i>Cyanophytes</i>	C16 :0	23.566 $\pm$ 2.069	17.601 $\pm$ 1.973	21.585 $\pm$ 3.337	11.392 $\pm$ 6.397	10.651 $\pm$ 7.958	12.005 $\pm$ 6.45	18.587 $\pm$ 9.179	11.459 $\pm$ 0.988	15.930 $\pm$ 0.987	19.071 $\pm$ 1.877	<0.01 (**)
	C16 :1n7	2.363 $\pm$ 0.607	1.678 $\pm$ 0.454	1.736 $\pm$ 1.002	1.353 $\pm$ 0.385	3.048 $\pm$ 1.584	2.932 $\pm$ 1.800	2.613 $\pm$ 0.763	4.603 $\pm$ 0.304	1.429 $\pm$ 0.436	2.561 $\pm$ 0.944	<0.01 (**)
<i>Phytoplankton</i>	C16 :4n3	7.910 $\pm$ 1.617	0.512 $\pm$ 0.161	0.477 $\pm$ 0.190	4.197 $\pm$ 3.356	7.744 $\pm$ 2.221	6.283 $\pm$ 4.326	4.116 $\pm$ 2.848	6.157 $\pm$ 0.592	1.523 $\pm$ 0.728	1.260 $\pm$ 0.331	<0.001 (***)
	C18 :0	5.358 $\pm$ 0.040	8.505 $\pm$ 1.087	7.555 $\pm$ 0.574	6.961 $\pm$ 3.514	9.011 $\pm$ 0.866	8.401 $\pm$ 4.036	8.403 $\pm$ 1.720	7.049 $\pm$ 0.095	4.748 $\pm$ 0.399	5.004 $\pm$ 0.172	<0.05 (*)
	C16 :4n3	7.910 $\pm$ 1.617	0.512 $\pm$ 0.161	0.477 $\pm$ 0.190	4.197 $\pm$ 3.356	7.744 $\pm$ 2.221	6.283 $\pm$ 4.326	4.116 $\pm$ 2.848	6.157 $\pm$ 0.592	1.523 $\pm$ 0.728	1.260 $\pm$ 0.331	<0.01 (**)
	C18 :2n6	1.983 $\pm$ 0.021	1.249 $\pm$ 0.290	1.765 $\pm$ 0.849	2.826 $\pm$ 1.519	2.865 $\pm$ 0.333	5.178 $\pm$ 2.281	3.507 $\pm$ 0.875	3.083 $\pm$ 0.400	35.268 $\pm$ 4.943	26.521 $\pm$ 3.229	<0.01 (**)

Values are expressed as means  $\pm$  SD of ten pooled clams for each site and three biological replicates were produced. Significant difference is given by asterisk at 0.05; \*p < 0.05; \*\*p < 0.01 and \*\*\*p < 0.001 using the ANOVA test (Tukey HSD).




FIGURE 5. Principal composition analysis (PCA) of trophic markers of *V. decussata* collected along the Tunisian Coast. (A) Correlation circle variables with the factorial axes of trophic markers and (B) individual projections on the factorial design of sampled sites (1×2).

Bizerte lagoon (S1), Chekly (S2), Baie (S3), Louza (S4), Zabbousa (S5), Boussaid (S6), Bunglow (S7), Maoumma (S8), Zarrat (S9) and Boughrara lagoon (S10).

green algae and urban discharge. PCA was applied to the fatty acid trophic marker proportions (Figure 5). A high % of variability accounted for the two first principal components (61.26%). Four indicator food sources of diatoms and dinoflagellates were found in all the *V. decussata* diets. However, green algae and phytoplankton markers characterized clams from all sampled sites except S2 and S3 ( $p < 0.01$ ). Almost all sites were characterized by cyanophyte diet sources, except for the Southern Tunisian sites (S9 and S10) ( $p < 0.01$ ). Two urban discharges and one bacteria fatty acid marker were found with highest proportion in the clams sampled from S9 and S10 ( $p < 0.001$ ). Detritus markers increased in the *V. decussata* tissues collected from S1, S2, S3, S4, S5, S6, S7 and S8 ( $p < 0.05$ ). However, the zooplankton marker characterized the diet of the clams sampled from S3 and S2 ( $p < 0.01$ ).

#### 4. DISCUSSION

In the present study, the PUFA content in *V. decussata* tissues was present at high values during the sampling period in all stations. PUFA were followed by SFA and then MUFA. These results are in agreement with previous studies carried out on the fatty acid composition of *V. decussata* from other Mediterranean Coasts (Ojea *et al.*, 2004) and of other bivalve species (Boussoufa *et al.*, 2011; Costa *et al.*, 2017) in which the authors showed a selective accumulation of PUFA among the fatty acid groups. MUFA were significantly lower in all stations, except in Boughrara (S10) and Zarrat (S9) where the clams showed similar SFA and MUFA proportions in their tissues. Changes in the PUFA

levels were inversely proportional to those of SFA. This result indicates that the unsaturation amount of the FA increased the water temperature was lower. The Pearson correlation test between the water temperature of different stations and the percentage of its n-3 and n-6 PUFA in each group of clams was applied. The results showed that water temperature was negatively correlated with (n-3) PUFA proportions ( $p < 0.003$ ;  $r = -0.4383$ ) and positively correlated with (n-6) PUFA proportions ( $p < 0.000$ ;  $r = 0.581$ ). This might be due to the greater levels of PUFA, which are necessary for maintaining the membrane fluidity of bivalves' cells during colder seasons (Irisarri *et al.*, 2014).

Regarding the FA profile, Louza (S4) clams contained significantly higher amounts of PUFA. However, the lagoon stations (S1, S2, S3 and S10) showed the lowest PUFA proportions. This can be explained by the fact that Louza (S4) was the least polluted station in our coast (Chalgmi *et al.*, 2014) compared to the lagoons, which were highly polluted (Bejaoui *et al.*, 2017) and showed degradation in their PUFA percentages. According to Di Salvatore *et al.*, (2013), a high proportion of PUFA reduces the vulnerability to lipid peroxidation and preserves the proper membrane changeability. Inversely, a reduction in PUFA levels could respond to oxidative damage which can be formed by heightened biotic and abiotic aspects in the ecosystem. The clams from other stations showed a similar distribution in their fatty acid signatures, therefore suggesting the use of similar food sources. In fact, the percent of PUFA recorded in our study was highly correlated with chlorophyll a and suspended matter. This confirms that changes in fatty acid composition are

strictly linked to available food and high levels of PUFA correspond with good nutritional conditions, as reported by Ojea *et al.*, (2004).

On the other hand, the fatty acid composition of *V. decussata* from the Tunisian Coast presented a prevalence of unsaturated fatty acids (UFA) throughout the study sites. According to many investigations, this higher amount of UFA was specific to a healthy marine species (Ghribi *et al.*, 2018). In fact, the PUFA levels in clam tissues reached their highest values, which coincided with an important amount of Chl a and lower T °C. Our results suggest that the unsaturation degree increased during the cold season, which maintained membrane fluidity as well as modulating gene expression and the principal keys of eicosanoids (Hochachka and Somero, 2002; Idayachandiran *et al.*, 2014). The change in fatty acid profile of the clams' tissues was influenced by T, which was probably due to the homeoviscous adaptation remodeling membrane lipids by changes in phospholipids, fatty acid and cholesterol contents, as reported previously by Hazel (1995). Our findings corroborate with previous studies carried out on bivalves (Irisarri *et al.*, 2014).

The multivariate analysis of the TFA data, including all samples showed a large difference among regions, explaining a clear north and south distinction along the Tunisian Coastline. Such spatial variation is most likely related to distinct trophic conditions (suspended materials, Chlorophyll a). The FCA analysis of the fatty acid composition was created by two prominent groups. The first was also formed by two sub-groups: The first sub-group included *V. decussata* sampled from the North of Tunisia (group A) (North and Bizerte Lagoons: S1, S2 and S3) characterized by a high proportion of some PUFA fatty acids, such as C22:6n-3. The second group's species were sampled from the center and north-south (group B) coastal waters. This sub-group was characterized by a lower percentage of dissimilarity (70%) than the first one. The FA composition of the clams sampled from S10 and S9 showed an independent group (group C) defined mostly by three polyunsaturated FA (C18:2n-6).

Bivalve species living in a shallow depth of the water column were directly influenced by the wide variety of potential food sources such as diatoms, dinoflagellates and zooplankton; and could have further complex trophic variation (Budge *et al.*, 2001). The fatty acid trophic markers used in this study revealed that diatoms and dinoflagellates were the primary incorporated food source of *V. decussata* from the Tunisian Coast. Those two diets are distributed in unequal levels over the sampling stations. In fact, the northern stations were dominated by a high proportion of diatom trophic markers such as sum C16 specific and C20:5n-3. Several studies carried out on some bivalve species showed that the level of C20:5n-3 was mainly derived from diatoms

(Budge *et al.*, 2001). However, the sum of C16:1n-7+C18:1 indicated that the base diet of diatoms was 2 times higher in the southern station (Zarrat and Boughrara lagoon) than in other areas. This unequal geographical dispersion of diatoms was confirmed by several studies showing the differential distribution of diatoms in the northern, central and southern part of Tunisian waters (Feki *et al.*, 2008; Chérif *et al.*, 2011). Concerning dinoflagellate markers, high levels of C16:0 and C18:4n-3 were observed in Bizerte, and the Northern and Southern Lagoons of Tunisia, which were significantly higher in the two southern stations (Gulf of Gabes). Like diatoms, numerous studies approved the geographical distribution of dinoflagellates among the Tunisian waters (Aissaoui *et al.*, 2012). However, the potential difference between diatoms and dinoflagellates was illustrated by the C22:6n-3/C20:5n-3 ratio (Dalsgaard *et al.*, 2003), which varied slightly among sampling sites and marked a significant high proportion in the Bizerte Lagoon. This result may be explained by the abundance of dinoflagellates and the prevalence of this diet compared to diatoms in this lagoon (Chérif *et al.*, 2011). These findings also suggest that the Bizerte Lagoon has a higher trophic position than the other stations. This position was explained by the high level of suspended matter and chlorophyll a signaled in this area and causing the eutrophication of this lagoon. Furthermore, water temperature, suspended matter, Chlorophyll a, currentology and especially wave amplitudes varied greatly among stations. These abiotic parameters have an important influence on the diet of *V. decussata* in these sites. Considering the difference between lagoon and sea, the wave amplitudes were higher in the coastal area of the Gulf of Gabes (Hattour and Ben Mustapha, 2013) than lagoons such as Bizerte, Boughrara and the North Lagoon of Tunisia. These later were characterized by low hydrodynamics causing water stagnation (Guetat *et al.*, 2012) and then eutrophication of these areas. This phenomenon could generate a physiological stress in bivalves reflected by a passive filter feeding and even an inhibition of the feeding process as a result of shell closure (Wildish and Kristmanson, 2005).

Similar studies carried on bivalves' feeding strategies showed the abundance of diatoms and dinoflagellates in their diet and a preference for dinoflagellates in colder down-welling seasons (Irisarri *et al.*, 2014). The diatom-dinoflagellate ratio was also used in several studies to distinguish among the diet in bivalves from different regions like Coral Sea, New Zealand and Tasmanian areas (Parrish *et al.*, 2015). In this context, Turki, (2004) signaled the presence of 61 species of dinoflagellates in the Bizerte Lagoon versus only 12 species of diatoms. Another typical trophic marker ratio C16:1/C16:0 used to deduce a main diatom versus a dinoflagellate diet (Auel *et al.*, 2002), was followed. In fact, the

results showed that the levels of C16:1/C16:0 ratios were lower than 1% in all sampling stations.

In the present study, the highest proportion of C18:2n-6 and C16:4n-3 in the soft tissue of *V. decussata* sampled from the Tunisian waters during winter affirmed the pervasiveness of phytoplankton as a food source. The results showed that the higher proportion of these two trophic markers was observed in clams from S1, S4, S5, S6, S7, S8, S9 and S10 compared to those from S2 and S3. Previous studies have shown that the same fatty acids can be used to suggest the phytoplankton integration of bivalves into wild seafood (Redmond *et al.*, 2010).

The Principal Component Analysis showed a geographical variation in the diet position of clams among sampling sites. Concerning the TFA composition of food sources, clams from S9 and S10 had significantly higher percentages of bacteria markers of (C16 and C18 PUFA), some green algae markers (C15:0 and 18:1n-7), and a lower proportion of diatoms (C20:5n-3) and dinoflagellate fatty acid markers (C22:6n-3) than from other stations. Further, the increase in bacterial fatty acid markers in the clams from S9 and S10 during the winter was accompanied by a high proportion of the two urban discharge (C18:1 and sum C18:1 + C18:2n-6). In fact, the Boughrara Lagoon (S10) and Zarrat (S9) are well known by their relatively high degrees of pollution (Rabaoui *et al.*, 2013) compared to other southern Tunisian waters. This fact could explain the development of the microbial loops and then their decomposition in these areas. In this context, Fernandez-Jover *et al.*, (2007) showed the usefulness of the determination of fatty acid trophic markers to identify the human impact on an animal's diet.

Herein, results showed a significant abundance of zooplankton in the diet of *V. decussata* sampled from the North Lagoon (S2, S3) as compared to other sampling areas. Our results are in agreement with those reported in several studies showing the abundance of zooplankton in the Gulf of Tunisia (Ben Lamine *et al.*, 2012). Concerning detritus FA indicators, the results showed the abundance of this diet in *V. decussata* sampled from the northern and central Tunisian waters. These specimens were characterized by high proportions of C22: n-6 PUFA and the sum (n-6 + n-3) HUFA compared to clams from other stations. Specimens of *V. decussata* from the north and the central stations (Bizerte, Chekly, Baie, Lousa and Zabbousa locations) were characterized by lower percentages of n-6 HUFA than n-3 HUFA. Clams from the southern Tunisian waters were dominated by n-6 HUFA compared to n-3 HUFA. Considering that n-6 HUFA was originated from terrestrial organic matter and n-3 HUFA was related to plankton detritus (Alfaro *et al.*, 2006), the clams from the north and center stations are considered to have an important energy source derived from planktonic detritus, compared to those from southern Tunisian waters.

## 5. CONCLUSIONS

This study revealed a significant geographical variation in the fatty acid composition of *V. decussata* related to environmental parameters of ten locations from the Tunisian Coast, thus revealing a good nutritional quality of the species marked by the abundance of PUFA especially DHA and EPA in all the sampled clams. To distinguish the feeding strategies and trophic links of the clam *V. decussata*, a fatty acid trophic marker analysis was conducted in clams from different sites. The results showed a spatial variation in food accessibility for *V. decussata* from the north to the south of the Tunisian coastline. These findings demonstrate that the FA signature can provide considerable information on the feeding strategy and the living condition of native *V. decussata* which can be used to discriminate among populations.

## ACKNOWLEDGMENTS

This work was supported by the Physiology and Aquatic Environment Laboratory, Faculty of Sciences, University of Tunis El Manar. We would like to thank all the team of this laboratory, especially Mr. Hssan Mejri for his technical assistance with the gas chromatography analysis.

## DISCLOSURE

The authors declare no conflict of interest.

## REFERENCES

- Aissaoui A, Turki S, Ben Hassine OK. 2012. Occurrence of harmful dinoflagellates in the Punic harbors of Carthage Gulf of Tunis and their correlations with the physico-chemical parameters. *Bull. Instit. Natl. Scien. Techn. Mer* **39**, 127–148. ISSN:0330-0080.
- Albergamo A, Rigano F, Purcaro G, Mauceri A, Fasulo S, Mondello L. 2016. Free fatty acid profiling of marine sentinels by nanoLC-ESI-MS for the assessment of environmental pollution effects. *Sci. Total Environ.* **571**, 955–962. <https://doi.org/10.1016/j.scitotenv.2016.07.082>
- Alfaro AC. 2006. Evidence of cannibalism and benthic-pelagic coupling within the life cycle of the mussel, *Perna canaliculus*. *J. Exp. Mar. Biol. Ecol.* **329**, 206–217. <https://doi.org/10.1016/j.jembe.2005.09.002>
- Aminot A, Chaussepied C. 1983. Manuel des analyses chimiques en milieu marin. In Brest CNEXO ed. Centre national d'exploitation des océans, France, 400p.
- Auel H, Harjes M, De Rocha R, Stübing D, Hagen W. 2002. Lipid biomarkers indicate different ecological niches and trophic relationships of the Arctic hyperiid amphipods *Themisto abyssorum* and *Themisto libellula*. *Pol. Bio.* **25**, 374–383.
- Bejaoui S, Boussoufa D, Tir M, Gharsallah-Houas I, Ghram A, El Cafsi M, Soudani N. 2017. DNA damage and oxidative stress in digestive gland of *Venerupis decussata*, collected from two contrasting habitats in the southern Tunisian coast: Biochemical and histopathological studies. *Cah. Bio. Mar.* **58**, 123–135. <https://doi.org/10.21411/CBM.A.133C71C1>
- Ben Lamine Y, Daly YKO, Daly YN. 2012. Planktonic copepod community in the neritic area south western part of Tunis



- bay influenced by Meliane river supplies (South western Mediterranean Sea). *Bull. Instit. National. Scien. Techno. Mer.* **39**, 141–147. ISSN 330–80.
- Boussoufa D, Ghazali N, Viciano E, Navarro JC, EL Cafsi M. 2011. Seasonal variation in condition and fatty acid composition of coquina clam, *Donax trunculus* (Linnaeus 1758) (Mollusca: Bivalvia) from the Tunisian coast. *Cah. Bio. Mar.* **52**, 47–56.
- Budge SM, Prrish CC, McKenzie CH. 2001. Fatty acid composition of phytoplankton, settling particulate matter and sediments at a sheltered bivalves aquaculture site. *Mar. Chem.* **76**, 285–303. [https://doi.org/10.1016/S0304-4203\(01\)00068-8](https://doi.org/10.1016/S0304-4203(01)00068-8)
- Budge SM, Iverson SJ, Koopman HN. 2006. Studying trophic ecology in marine ecosystems using fatty acids: a primer on analysis and interpretation. *Mar. Mamm. Sci.* **22**, 759–801. <https://doi.org/10.1111/j.1748-7692.2006.00079.x>
- Capillo G, Silvestro S, Sanfilippo M, Fiorino E, Giangrosso G, Ferrantelli V, Vazzana I, Faggio C. 2018. Assessment of Electrolytes and Metals Profile of the Faro Lake (Capo Peloro Lagoon, Sicily, Italy) and Its Impact on *Mytilus galloprovincialis*. *Chem. Biodivers.* **15**, e1800 D44. <https://doi.org/10.1002/cbdv.201800044>
- Chalghmi H, Bourdineaud JP, Haouas Z, Gourves PY, Zrafi I, Saidane-Mosbah D. 2014. Transcriptomic, Biochemical, and Histopathological Responses of the Clam *Ruditapes decussatus* from a Metal-Contaminated Tunis Lagoon. *Arch. Environ. Contam. Toxicol.* **70**, 241–56. <https://doi.org/10.1007/s00244-015-0185-0>
- Cecchi G, Basini S, Castano C. 1985. Méthanalyse rapide des huiles en solvant. *R. Franç. Corps Gras* **4**.
- Chérif W, El Bour M, Daly YKO. 2011. Screening de l'activité anti microfouling d'algues verts récoltées sur la côte nord Tunisienne. *Bull. Instit. Nat. Sci. Techn. Mer.* **38**, 131–138. ISSN 0330-0080.
- Costa R, Alberghamo A, Piparo M, Zaccane G, Capillo G, Manganaro A, Mondello L. 2017. Multidimensional gas chromatographic techniques applied to the analysis of lipids from wild-caught and farmed marine species. *Eur. J. Lipid. Sci. Technol.* **119**, 1600043. <https://doi.org/10.1002/ejlt.201600043>
- Dalsgaard J, John M, Kattner G, Muller-Navarra D, Hagen W. 2003. Fatty acid trophic markers in the pelagic marine environment. *Adv. Mar. Biol.* **46**, 225–340. [https://doi.org/10.1016/S0065-2881\(03\)46005-7](https://doi.org/10.1016/S0065-2881(03)46005-7)
- Desvillettes CH, Bourdier G, Amblard CH, Barth B. 1997a. Use of fatty acids for the assessment of zooplankton grazing on bacteria, protozoans and microalgae. *Fresh. Water. Biol.* **38**, 629–637. <https://doi.org/10.1046/j.1365-2427.1997.00241.x>
- Fanelli E, Papiol V, Cartes JE, Rumolo P, Lopez-Pérez C. 2013. Trophic webs of deep megafauna on mainland and insular slopes of the NW Mediterranean: a comparison by stable isotope analysis. *Mar. Ecol. Prog. Ser.* **490**, 199–221. <https://doi.org/10.3354/meps10430>
- Fernandez-Jover D, Jimenez JAL, Sanchez-Jerez P, Bayle-Sempere J, Casaldueiro FG, Lopez FJM, Dempster T. 2007. Changes in body condition and fatty acid composition of wild Mediterranean horse mackerel (*Trachurus mediterraneus*, Steindachner, 1868) associated to sea cage fish farms. *Mar. Environ. Res.* **63**, 1–18. <https://doi.org/10.1016/j.marenvres.2006.05.002>
- Feki W, Hamza A, Bel Hassen M, Rebai A. 2008. Les efflorescences phytoplanctoniques dans le golfe de gabes (Tunisie) au cours de dix ans de surveillance (1995-2005). Institut National des Sciences et Technologies de la Mer **35**, 105–116.
- Folch J, Lees M, Stanley GHS. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* **226**, 497–509.
- Gomes T, Alberghamo A, Costa R, Mondello L, Dugo G. 2017. Potential use of proteomics in shellfish aquaculture: from assessment of environmental toxicity to evaluation of seafood quality and safety. *Cur. Org. Chem.* **21**, 402–425. <https://doi.org/10.2174/1385272820666161102121232>
- Ghribi F, Boussoufa D, Aouini F, Bejaoui S, Chetoui I, Rabeh I, El Cafsi M. 2018. Seasonal variation of biochemical composition of Noah's ark shells (*Arca noae* L. 1758) in a Tunisian coastal lagoon in relation to its reproductive cycle and environmental conditions. *Aquat. Living Resour.* <https://doi.org/10.1051/alr/2018002>
- Guétat F, Sellem F, Akrouf F, Brahim M, Atoui A. 2012. Etat environnemental de la lagune de Boughrara et ses alentours deux ans après les travaux d'aménagement et d'élargissement du pont d'El Kantra. *Bull. Inst. Nat. Scien. Tech. Mer de Salammbô*, **v39**. ISSN: 0330-0080.
- Hamida L, Medhioub M, Cochard JC, Romdhane M, Le Pennec M. 2004. A comparative study of the reproductive cycle of *Ruditapes decussatus* under natural (South of Tunisia) and controlled conditions (hatchery). *Cah. Biol. Mar.* **45**, 291–303.
- Hattour A, Ben Mustapha A. 2013. Le couvert végétal marin du golfe de gabes: cartographie et réseau de surveillance de l'herbier de posidonie. *Bull. Instit. Nat. Sci. Techn. Mer.* **164** pp.
- Hazel JR. 1995. Thermal adaptation in biological membranes: is homeoviscous adaptation the explanation? *Annu. Rev. Physiol.* **57**, 19–42.
- Hochachka PM, Somero GN. Biochemical adaptation. Mechanism and process in physiological evolution, Princeton University Press, Oxford, 2002, 480 p.
- Idayachandiran G, Muthukumar A, Kumaresan S, Balasubramanian T. 2014. Nutritional Value of Marine Bivalve, *Donax cuneatus* 86 (Linnaeus, 1758) from Cuddalore Coastal Waters, Southeast Coast of India. *Inventi Impact Life Style* **1**, 15–19.
- Irisarri J, Fernandez-Reiriz MJ, Cranford PJ, Labarta U. 2014. Effects of seasonal variation in phytoplankton on the bio-energetic responses of mussels (*Mytilus galloprovincialis*) held on a raft in the proximity of sea bream (*Pagellus bogaraveo*) net-pens. *Aqua.* **428–429**, 41–53. <https://doi.org/10.1016/j.aquaculture.2014.02.030>
- Kharlamenko IV, Wurzburg L, Peters J, Borisovets EE. 2015. Fatty acid compositions and trophic relationships of shelled molluscs from the kuril-Kamchatka Trench and the adjacent abyssal plain. *Deep-sea Res. II* **111**, 389–398. <https://doi.org/10.1016/j.dsr2.2014.09.002>
- Leveillé JC, Amblard C, Bourdier G. 1997. Fatty acids as specific algal markers in a natural lacustrine phytoplankton. *J. Plankton. Res.* **19**, 469–490. <https://doi.org/10.1093/plankt/19.4.469>
- Li R, Watanabe MM. 2001. Fatty acid profiles and their chemotaxonomy in planktonic species of *Anabaena* (Cyanobacteria) with straight trichomes. *Phytoch.* **57**, 727–731. [https://doi.org/10.1016/S0031-9422\(01\)00082-6](https://doi.org/10.1016/S0031-9422(01)00082-6)
- McLusky SD, Elliott M. 2004. The Estuarine Ecosystem: Ecology, threats and management. University press Third Edition, Oxford.
- Natalotto A, Sureda A, Maisano M, Spanò N, Mauceri A, Deudero S. 2015. Biomarkers of environmental stress in gills of *Pinna nobilis* (Linnaeus 1758) from Balearic Island. *Ecotoxicol. Environ. Saf.* **122**, 9–16. <https://doi.org/10.1016/j.ecoenv.2015.06.035>
- Ojea J, Pazos AJ, Martinez D, Novoa S, Sanchez JL, Abad M. 2004. Seasonal variation in weight and biochemical composition of the tissues of *Ruditapes decussatus* in relation to the gametogenic cycle. *Aqua.* **238**, 451–468. <https://doi.org/10.1016/j.aquaculture.2004.05.022>
- Pagano M, Capillo G, Sanfilippo M, Palato S, Trischitta F, Manganaro A, Faggio C. 2016. Evaluation of Functionality and Biological Responses of *Mytilus galloprovincialis* after Exposure to Quaternium-15 (Methenamine 3-Chloroallylchloride). *Molecules* **21**, 144. <https://doi.org/10.3390/molecules21020144>
- Parrish CC, Pethybridge H, Young JW, Nichols PD. 2015. Spatial variation in fatty acid trophic markers in albacore tuna from the southwestern Pacific Ocean-A potential 'tropicalization' signal. *Deep. sea. Res. II* **113**, 19–207. <https://doi.org/10.1016/j.dsr2.2013.12.003>
- Pernet F, Malet N, Pastoureaud A, Vaquer A, Quéré C, Dubroca L. 2012. Marine diatoms sustain growth of bivalves in a Mediterranean lagoon. *J. Sea. Res.* **68**, 20–32. <https://doi.org/10.1016/j.seares.2011.11.004>

- Prato E, Danieli A, Maffia M, Biantolino F. 2010. Lipid and Fatty Acid Compositions of *Mytilus galloprovincialis* Cultured in the Mar Grande of Taranto (Southern Italy): Feeding Strategies and Trophic Relationships. *Zool. Stu.* **49**, 211–219.
- Rabaoui L, Balti R, El Zrelli R, Tlig-Zouari S. 2013. Assessment of heavy metal pollution in the gulf of Gabes (Tunisia) using four mollusc species. *Med. Mar. Sci.* **15**. <https://doi.org/10.12681/mms.504>
- Redmond KJ, Magnesen T, Hansen PK, Strand O, Meier S. 2010. Stable isotopes and fatty acids as tracers of the assimilation of salmon fish feed in blue mussels (*Mytilus edulis*). *Aqua.* **298**, 202–210. <https://doi.org/10.1016/j.aquaculture.2009.10.002>
- Rocchetta I, Pasquevich MY, Heras H, De Molina MDCR, Luquet CM. 2014. Effects of sewage discharges on lipid and fatty acid composition of the Patagonian bivalve *Diplodon chilensis*. *Mar. Poll. Bull.* **79**, 211–219. <https://doi.org/10.1016/j.marpolbul.2013.12.011>
- Sakdullah A, Tsuchiya M. 2009. The origin of particulate organic matter and the diet of tilapia from an estuarine ecosystem subjected to domestic wastewater discharge: fatty acid analysis approach. *Aqua. Ecol.* **43**, 577–589. <https://doi.org/10.1007/s10452-008-9195-6>
- Salvatore P di, Calcagno JA, Ortiz N, De Molina MDCR, Sabatini SE. 2013. Effect of seasonality on oxidative stress responses and metal accumulation in soft tissues of *Aulacomya atra*, a mussel from the south Atlantic Patagonian coast. *Mar. Environ. Res.* **92**, 244–252. <https://doi.org/10.1016/j.marenvres.2013.10.004>
- Turki S. 2004. Suivi des microalgues planctoniques toxiques dans les zones de production, d'élevage des mollusques bivalves et d'exploitation des oursins du nord de la Tunisie. *Bull. Instit. Nat. Sci. Techn. Mer.* **31**, 83–96.
- Wang S, Jin B, Qin H, Sheng Q, Wu J. 2015. Trophic Dynamics of Filter Feeding Bivalves in the Yangtze Estuarine Intertidal Marsh: Stable Isotope and Fatty Acid Analyses. *PLOS* **10**, e0135604. <https://doi.org/10.1371/journal.pone.0135604>
- Wildish D, Kristmanson D. 2005. Benthic suspension feeders and flow. Cambridge University, New York.